

RUBINSTEIN DISTANCE ON CONFIGURATIONS SPACES

L. DECREUSEFOND AND N. SAVY

ABSTRACT. By a method inspired of the Stein's method, we derive an upper-bound of the Rubinstein distance between two absolutely continuous probability measures on configurations space. As an application, we show that the best way to approximate a Modulated Poisson Process (see below for the definition) by a Poisson process is to equate their intensity.

1. INTRODUCTION

According to the Kantorovitch approach, the optimal transportation problem or Monge-Kantorovitch problem (MKP for short) reads as follows: given two probability measures μ and ν on a Polish space X and a cost function c on $X \times X$, does there exist a probability measure γ on $X \times X$ which minimizes $\int c \, d\beta$ among all probability measures β on $X \times X$ with first (respectively second) marginal μ (respectively ν) ? The first step is to determine whether or not there exists a probability measure γ such that $\int c \, d\gamma$ is finite. In the solved cases, a few criterion are known. The oldest one (see [17]), for quadratic cost, stands that such a measure exists provided that μ and ν have finite second moments. Still for the quadratic cost, if μ is a Gaussian measure on a finite or infinite dimensional space and $\nu = L\mu$, then the distance is finite whenever L has a finite Boltzmann entropy [9]. In the same reference, a bound may also be found for the Rubinstein distance, i.e., when c is a distance function. We are interested in the evaluation of the Rubinstein distance on configurations space, i.e., for locally finite point processes. The first point to be stressed is that we have several reasonable distance between configurations. To name only the two we will investigate here: there is the total variation distance when configurations are viewed as atomic measures, and there also is a distance with a greater geometric flavor, which is defined in (8). To these two distances correspond two different notions of Lipschitz continuous functions. This is of some great importance since the Kantorovitch-Rubinstein duality allows us to write the Rubinstein distance as a maximization problem on the set of Lipschitz functions, see (1). Moreover, it is well known in finite dimension that Lipschitz functions are “almost” differentiable and that their differential is bounded. It turns out that the two usual gradients introduced on configurations space (see [1],[16]) are the good tools to obtain an analog result on configurations space. Once we have a gradient, we usually introduce the divergence (as its adjoint) and then a number operator (as the composition of the divergence and the gradient), hence an Ornstein-Uhlenbeck (see [4]) semi-group. At this point, it is useful to invoke the Stein's method which is a very efficient tool to obtain stochastic bounds notably for point processes (see [6, 5, 21, 2]). In essence, Stein's method compares the expectations of two distinct random variables by “embedding” them in the evolution of an ergodic Markov process and by then looking

Date: February 1, 2008.

Key words and phrases. Configurations space, Malliavin calculus, Monge-Kantorovitch problem, Point processes, Rubinstein distance, Stein's method.

backward at the evolution of this process, from infinity to time 0. With very sketchy notations, imagine that we have two smooth functions α and β with the same limit at infinity, then $\alpha(0) - \beta(0)$ may be evaluated by computing

$$\int_0^\infty |\alpha'(s) - \beta'(s)| \, ds.$$

Controlling the difference of derivatives yields to a bound on the difference at time 0. This is exactly the principle at work in Theorem 1. This method reminds of the so-called semi-group method often used in proofs of concentration inequality [13]. In Section 2, we present the generic principle of our method and we apply it to the different distances on configurations spaces in Section 3. Section 4 is devoted to the application of these results on the approximation of Markov modulated Poisson Process (MMPP) by a Poisson process. The motivation for this part comes from queueing theory where MMPP are widely used because of their versatility, useful to model a wide range of real systems [11, 14, 15] and of the persistence of their Markovianity. Unfortunately, these processes are affected by the curse of dimensionality: it is often the case that we must invert linear systems with a so huge number of variables it becomes unfeasible. It is thus of crucial importance to reduce the cardinality of state space. The extreme situation is when this space is reduced to one point, i.e., when an MMPP is a Poisson process. We find by the method developed in the beginning of this paper, a bound on the Rubinstein distance between an MMPP and a Poisson process. Optimizing this bound yields to the well known rule of thumb which consists in taking as the optimal intensity, the average intensity of the MMPP. Our result is not then astoundingly original but it shows that by proceeding along this line, we can control the error for any functional such as loss probability or others.

2. GENERIC SCHEME

Let X be a Polish space and d a lower-semi-continuous distance function on $X \times X$, which does not necessarily generate the topology on X . We will denote by $d - \text{Lip}_m$ the set of Lipschitz continuous F from X to $\mathbb{R}$ with Lipschitz constant m :

$$|F(x) - F(y)| \leq m d(x, y),$$

for any $(x, y) \in X^2$. For two probability measures μ and ν on X , the optimal transportation problem associated to d consists in evaluating

$$\mathcal{T}_d(\mu, \nu) = \inf_{\gamma \in \Sigma(\mu, \nu)} \int_{X \times X} d(x, y) \, d\gamma(x, y),$$

where $\Sigma(\mu, \nu)$ is the set of probability measures on $X \times X$ with first (respectively second) marginal μ (respectively ν). According to [8, 20], this minimum is equal to

$$(1) \quad \mathcal{T}_d(\mu, \nu) = \sup_{F \in d - \text{Lip}_1} \int F \, d(\mu - \nu).$$

We consider the situation where $X = \Gamma_\Lambda$ is the configurations space on a Lusin space Λ , i.e.,

$$\Gamma_\Lambda = \{\eta \subset \Lambda; \eta \cap K \text{ is a finite set for every compact } K \subset \Lambda\}.$$

We identify $\eta \in \Gamma_\Lambda$ and the positive Radon measure $\sum_{x \in \eta} \varepsilon_x$. Throughout this paper, Γ_Λ is endowed with the vague topology, i.e., the weakest topology such that for all $f \in \mathcal{C}_0$ (continuous with compact support on Λ), the maps

$$\eta \mapsto \int_\Lambda f \, d\eta = \sum_{x \in \eta} f(x)$$

are continuous. When f is the indicator function of a subset B , we will use the shorter notation $\eta(B)$ to denote the integral of $\mathbf{1}_B$ with respect to η . We denote by $\mathcal{B}(\Gamma_\Lambda)$ the corresponding Borel σ -algebra. The probability space under consideration will then be $(\Gamma_\Lambda, \mathcal{B}(\Gamma_\Lambda), \mu)$. We need some additional structure.

Hypothesis 1. Assume now that we have :

- a kernel Q on $X \times \Lambda$, i.e., such that $Q(A, \cdot)$ is measurable as function on Λ for any $A \in \mathcal{B}(\Gamma_\Lambda)$ and $Q(\cdot, s)$ is a σ -finite measure on X for any $s \in \Lambda$,
- a map ∇ , defined on a subset $\text{Dom } \nabla$ of $L^2(\mu)$, such that, for any $F \in \text{Dom } \nabla$,

$$\int_X \int_\Lambda |\nabla_s F|^2 Q(\omega, ds) d\mu(\omega) < +\infty.$$

We say that a process $u(\omega, s)$ belongs to $\text{Dom } \delta$ whenever, there exists a constant c independent of F such that for any $F \in \text{Dom } \nabla$,

$$\left| \mathbb{E} \left[\int_\Lambda \nabla_s F u(s) Q(\omega, ds) \right] \right| \leq c \|F\|_{L^2(\Omega)}.$$

For such a process u , we define δu by

$$(2) \quad \int_X \int_\Lambda \nabla_s F(\omega) u(\omega, s) Q(\omega, ds) d\mu(\omega) = \int_X F \delta u d\mu.$$

Definition 1. We say that (∇, Q) has the Rademacher property whenever $F \in d - \text{Lip}_1$ implies $\nabla F \in \text{Dom } \nabla$ and

$$(3) \quad |\nabla_s F| \leq 1, \quad Q(\omega, \cdot) d\mu\text{-almost-surely.}$$

Consider for $F \in \text{Dom } \nabla$, the (formal) equation

$$(4) \quad \frac{d}{dt} X_t = -\delta \nabla X_t, \quad X_0 = F.$$

If this equation has one and only one solution for each $F \in \text{Dom } \nabla$, we then have a μ -self-adjoint semi-group $(P_t, t \geq 0)$, usually called the Ornstein-Uhlenbeck semi-group: $P_t F = X_t$ where X_t is the solution of (4).

Definition 2. The Ornstein-Uhlenbeck is said to be ergodic whenever $\lim_{t \rightarrow +\infty} P_t F = \int_X F d\mu$.

Theorem 1. Assume that hypothesis 1 holds. Let ν be another probability measure on X absolutely continuous with respect to μ . We denote by L the Radon-Nikodym derivative of ν with respect to μ . If (∇, Q) has the Rademacher property and if the Ornstein-Uhlenbeck semi-group is ergodic then:

$$(5) \quad \mathcal{T}_d(\mu, \nu) \leq \int_{X \times \Lambda} \int_0^{+\infty} |\nabla_s P_t L| dt Q(\omega, ds) d\mu(\omega).$$

Proof. According to the fundamental Lemma of analysis,

$$\begin{aligned} \int_X F d\mu - F &= \int_0^{+\infty} \frac{d}{dt} P_t F dt, \\ &= \int_0^{+\infty} \delta \nabla P_t F dt, \\ &= \int_0^{+\infty} P_t \delta \nabla F dt. \end{aligned}$$

Since $d\nu/d\mu = L$,

$$\begin{aligned}
\int_X F \, d\mu - \int_X F \, d\nu &= \int_X \left(\int_X F \, d\mu - F \right) d\nu \\
&= \int_X \left(\int_0^{+\infty} P_t \delta \nabla F \, dt \right) d\nu \\
&= \int_X \int_0^{+\infty} P_t \delta \nabla F L \, dt \, d\mu \\
&= \int_{X \times \Lambda} \int_0^{+\infty} \nabla_s F \, \nabla_s P_t L \, dt \, Q(\omega, ds) \, d\mu(\omega).
\end{aligned}$$

Since (∇, Q) has the Rademacher property, we have an $L^\infty - L^1$ bound which yields to (5). $\square$

3. INSTANTIATIONS ON POISSON SPACE

Let ρ be a σ -finite measure on Λ and assume that μ is the Poisson measure of intensity ρ , i.e., the probability measure on Γ_Λ fully characterized by

$$\mathbb{E} \left[\exp \left(\int_\Lambda f \, d\eta \right) \right] = \exp \left(\int_\Lambda \left(e^{f(s)} - 1 \right) d\rho(s) \right).$$

3.1. Discrete gradient on Poisson space. For $F : X \rightarrow \mathbb{R}$, the discrete gradient of F , denoted by $\nabla^\sharp F$, is defined by

$$\nabla_s^\sharp F(\eta) = F(\eta + \varepsilon_s) - F(\eta).$$

We set $Q(\omega, ds) = d\rho(s)$ so that $\text{Dom } \nabla^\sharp$ is defined as the set of functionals such that

$$\mathbb{E} \left[\int_\Lambda |\nabla_s^\sharp F|^2 d\rho(s) \right] < +\infty.$$

We denote by $\delta^\sharp$ its adjoint in the sense of (2). The n -th iterated integral of a symmetric function f from Λ^n to $\mathbb{R}$ is defined as

$$J_n(f) = n! \int \dots \int_{0 \leq s_1 < s_2 < \dots < s_n} f(s_1, \dots, s_n) \, d(\eta - \rho)(s_1) \dots d(\eta - \rho)(s_n).$$

For a general function f ,

$$J_n(f) = \sum_{\sigma \in \mathcal{S}_n} \int \dots \int_{0 \leq s_1 < s_2 < \dots < s_n} f(s_{\sigma(1)}, \dots, s_{\sigma(n)}) \, d(\eta - \rho)(s_1) \dots d(\eta - \rho)(s_n).$$

It is well known [19, 16] that any square integrable functional on Γ_Λ can be written as

$$F = \sum_{n=0}^{+\infty} J_n(f_n),$$

where for any integer n , f_n is symmetric and belongs to $L^2(\rho^{\otimes(n)})$ and that

$$\nabla_s^\sharp F(\eta) = \sum_{n=1}^{+\infty} n J_{n-1}(f_n(\cdot, s)).$$

Moreover, the Ornstein-Uhlenbeck semi-group operates on chaos as:

$$(6) \quad P_t^\sharp F = \sum_{n=1}^{+\infty} e^{-nt} J_n(f_n)$$

From (6) and by dominated convergence, it is then easily seen that $P^\sharp$ is ergodic. We now choose the total variation as the distance of interest on Γ_Λ , i.e.,

$$d_1(\eta, \omega) = 2 \sup_{A \in \Lambda} |\eta(A) - \omega(A)|$$

Lemma 1. *For the distance d_1 on X , $(\nabla^\sharp, Q)$ has the Rademacher property.*

Proof. Consider $F \in d_1 - \text{Lip}_1$, by the very definition of the gradient:

$$\begin{aligned} |\nabla_s^\sharp F(\eta)| &\leq |F(\eta + \varepsilon_s) - F(\eta)| \\ &\leq d_1(\eta + \varepsilon_s, \eta) = 1. \end{aligned}$$

In the converse direction, consider ω and η be two locally finite but not finite configurations. If $d_1(\omega, \eta) = +\infty$, there is nothing to prove. If $d_1(\omega, \eta)$ is finite, $\omega\Delta\eta$ and $\eta\Delta\omega$ are finite, where $\omega\Delta\eta = \omega \setminus (\omega \cap \eta)$. Since $|\nabla_s^\sharp F(\eta)| \leq 1$, we get:

$$\begin{aligned} |F(\eta) - F(\omega)| &\leq |F(\eta \cap \omega \cup \eta\Delta\omega) - F(\eta \cap \omega)| + |F(\eta \cap \omega \cup \omega\Delta\eta) - F(\eta \cap \omega)| \\ &\leq (\eta\Delta\omega)(\Lambda) + (\omega\Delta\eta)(\Lambda) \\ &\leq 2 \max((\eta\Delta\omega)(\Lambda), (\omega\Delta\eta)(\Lambda)) \\ &= d_1(\eta, \omega). \end{aligned}$$

The Rademacher property is then established for $(\nabla^\sharp, Q)$. $\square$

Theorem 2. *Let μ and ν two probability measures on Γ_Λ such that $d\nu = L d\mu$. We have,*

$$\mathcal{T}_{d_1}(\mu, \nu) \leq \mathbb{E} \left[\int_\Lambda |(\text{Id} + \mathcal{L}^\sharp)^{-1} \nabla_s^\sharp L| d\rho(s) \right],$$

where $\mathcal{L}^\sharp = \delta^\sharp \nabla^\sharp$.

Proof. It is easily seen using chaos decomposition that

$$\nabla_s^\sharp P_t^\sharp F = e^{-t} P_t^\sharp \nabla_s^\sharp F \quad \text{for all } s \in \lambda, \text{ for all } t \in \mathbb{R}^+$$

and it is a general property of semi-groups and their generator that

$$\int_0^{+\infty} e^{-t} P_t^\sharp F dt = (\text{Id} + \mathcal{L}^\sharp)^{-1} F,$$

for any $F : \Omega \rightarrow \mathbb{R}$. We then infer from (5) that

$$\begin{aligned} \mathcal{T}_{d_1}(\mu, \nu) &\leq \mathbb{E} \left[\int_\Lambda \left| \int_0^{+\infty} e^{-t} P_t^\sharp \nabla_s^\sharp L dt \right| d\rho(s) \right] \\ &= \mathbb{E} \left[\int_\Lambda |(\text{Id} + \mathcal{L}^\sharp)^{-1} (\nabla_s^\sharp L)| d\rho(s) \right] \end{aligned}$$

$\square$

Remark 1. *Note that the very analog of this inequality on Wiener space was proved by a different though related way in [10].*

3.2. Derivation on Poisson space. In this section we introduce another stochastic gradient on Γ_Λ which is a derivation – see [1]. Let $V(\Lambda)$ be the set of $\mathcal{C}^\infty$ vector fields on Λ and $V_0(\Lambda) \subset V(\Lambda)$, the subset consisting of all vector fields with compact support. For $v \in V_0(\Lambda)$, for any $x \in \Lambda$, the curve

$$t \mapsto \mathcal{V}_t^v(x) \in \Lambda$$

is defined as the solution of the following Cauchy problem

$$(7) \quad \begin{cases} \frac{d}{dt} \mathcal{V}_t^v(x) &= v(\mathcal{V}_t^v(x)), \\ \mathcal{V}_0^v(x) &= x. \end{cases}$$

The associated flow $(\mathcal{V}_t^v, t \in \mathbb{R})$ induces a curve $(\mathcal{V}_t^v)^* \eta = \eta \circ (\mathcal{V}_t^v)^{-1}$, $t \in \mathbb{R}$, on Γ_Λ : If $\eta = \sum_{x \in \eta} \varepsilon_x$ then $(\mathcal{V}_t^v)^* \eta = \sum_{x \in \eta} \varepsilon_{\mathcal{V}_t^v(x)}$. We are then in position to define the notion of differentiability on Γ_Λ . A measurable function $F : \Gamma_\Lambda \rightarrow \mathbb{R}$ is said to be differentiable if for any $v \in V_0(\Lambda)$, the following limit exists:

$$\lim_{t \rightarrow 0} t^{-1} (F(\mathcal{V}_t^v(\eta)) - F(\eta)).$$

We then denote $\nabla_v^c F(\eta)$ the preceding limit. We denote by $\nabla_s^c F$ we corresponding gradient. It verifies :

$$\nabla_v^c F(\omega) = \int_\Lambda \nabla_s^c F(\omega) v(s) \, d\omega(s).$$

The square norm of $\nabla^c F$ is given by

$$\int_\Lambda \nabla_s^c F \, d\omega(s),$$

so that we are in the framework of Hypothesis 1 if we take

$$Q(\omega, ds) = d\omega(s) = \sum_{x \in \omega} \varepsilon_x(ds),$$

where ε_a is the Dirac mass in a . For a random variable $F : \Gamma_\Lambda \rightarrow \mathbb{R}$, and random process $u : \Gamma_\Lambda \times \Lambda \rightarrow \mathbb{R}$, we define the adjoint operator of ∇^c denoted by δ^c by:

$$\mathbb{E} [\langle \nabla^c F, u \rangle_{L^2(\rho)}] = \mathbb{E} [F \delta^2 u],$$

provided both sides exist, i.e.,

$$|\mathbb{E} [\langle \nabla^c F, u \rangle_{L^2(\rho)}]| \leq c \|F\|_{L^2}^2.$$

Consider now $\mathcal{L}^c = \delta^c \nabla^c$ and the associated semi-group semigroup $\{P_t^c, t \in \mathbb{R}\}$. The distance of interest is here the Wassertein's distance (see [7, 18]):

$$(8) \quad d_2(\eta_1, \eta_2) = \left[\inf \left\{ \int d_0(x, y) d\beta(x, y), \beta \in \Gamma_{\eta_1, \eta_2} \right\} \right]^{1/2},$$

where Γ_{η_1, η_2} denotes the set of $\beta \in \Gamma_{\Lambda \times \Lambda}$ having marginals η_1 and η_2 . The ergodicity of P^c is proved in [1] and the Rademacher property is the object of [18].

On the other hand, there is no known commutation relationships between ∇^c and P_t^c hence theorem 1 entails that

Theorem 3. *Let μ and ν two probability measures on Γ_Λ such that $d\nu = L \, d\mu$. We have,*

$$\mathcal{I}_{d_2}(\mu, \nu) \leq \mathbb{E} \left[\int_0^{+\infty} |\nabla^c P_t^c L| \, dt \right].$$

4. APPLICATIONS

4.1. Distance between two Poisson processes.

Theorem 4. *Consider μ and ν two Poisson probability measures on $\Lambda \subset \mathbb{R}^n$. The intensity of μ is the Lebesgue measure on Λ and that of ν is $h(s)ds$ with $h \in L^2(\Lambda)$ deterministic. Then, we have the following bound :*

$$(9) \quad \mathcal{T}_{d_1}(\mu, \nu) \leq C \|h\|_{L^1(\Lambda)} \exp\left(\frac{1}{2} \|h - 1\|_{L^2(\Lambda)}^2\right).$$

Proof. According to [12],

$$L(\omega) = \frac{d\nu}{d\mu}(\omega) = \exp\left(\int_{\Lambda} \ln h(s) d\omega(s) - \int_{\Lambda} (h(s) - 1) ds\right).$$

By definition,

$$\nabla_s^\# L(\omega) = L(\omega + \varepsilon_s) - L(\omega) = h(s)L.$$

Hence, according to Theorem 2,

$$\mathcal{T}_{d_1}(\mu, \nu) \leq C \|h\|_{L^1(\Lambda)} \mathbb{E} [|(I + \mathcal{L}^\#)^{-1} L|].$$

It is then well known, using for instance the chaos decomposition that

$$\mathbb{E} [|(I + \mathcal{L}^\#)^{-1} L|] \leq \mathbb{E} [L^2]^{1/2}$$

and

$$\begin{aligned} \mathbb{E} [L^2] &= \mathbb{E} \left[\exp\left(2 \int_{\Lambda} \ln h(s) d\omega(s) - 2 \int_{\Lambda} (h(s) - 1) ds\right) \right] \\ &= \mathbb{E} \left[\exp\left(\int_{\Lambda} \ln h^2(s) d\omega(s) - \int_{\Lambda} (h^2(s) - 1) ds\right) \right] \exp\left(\int_{\Lambda} (h(s) - 1)^2 ds\right) \\ &= \exp\left(\int_{\Lambda} (h(s) - 1)^2 ds\right). \end{aligned}$$

The proof is thus complete. $\square$

4.2. Distance between a Poisson process and a Markov modulated Poisson process. In this section we calculate a bound for the distance between a Poisson Process and an Markov modulated Poisson process (MMPP for short).

Definition 3. *Consider J an irreducible continuous time Markov chain with finite state space. We denote by m_J the finite number of states of J , Q_J the infinitesimal generator of J , π_J the stationary vector of Q_J . We assume also that we are given $(\lambda_1, \dots, \lambda_{m_J})$, m_J non-negative real numbers. We denote by Ψ the map which sends $i \in \{1, \dots, m_J\}$ to λ_i . An MMPP(J, Ψ) is a point process the intensity of which is given by $\Psi(J_s)ds$. This means that when J is in state (called a phase) k ($1 \leq k \leq m_J$) then the arrivals occurs according to a Poisson process of rate λ_k .*

A detailed description of the MMPP with an emphasis on applicability to modeling is given in [14] and references therein.

Theorem 5. *Let μ be a Poisson process of intensity λ and ν be a MMPP(J, Ψ) then for a given $T \in \mathbb{R}^+$,*

$$(10) \quad \mathcal{T}_{d_1}(\mu_T, \nu_T) \leq \mathbb{E} \left[\int_0^T \left| \frac{\Psi(J_s)}{\lambda} \right| \lambda ds \exp \left[\frac{1}{2} \int_0^T \left| \frac{\Psi(J_s)}{\lambda} - 1 \right|^2 \lambda ds \right] \right]$$

Proof. By the Girsanov formula we have:

$$L_T(\omega) = \frac{d\nu_T}{d\mu_T}(\omega) = \exp\left(\int_0^T \ln \frac{\Psi(J_s)}{\lambda} d\omega(s) - \int_0^T \left(\frac{\Psi(J_s)}{\lambda} - 1\right) \lambda ds\right).$$

Consider $\mathfrak{F}_J^T = \sigma\{J_s, s \leq T\}$, the history of J up to time T . Now,

$$\begin{aligned} \mathcal{T}_{d_1}(\mu_T, \nu_T) &= \sup_{F \in d_1\text{-Lip}_1} (\mathbb{E}_{\mu_T} [F(L_T - 1)]) \\ &= \sup_{F \in d_1\text{-Lip}_1} (\mathbb{E}_{\mu_T} [\mathbb{E}[F(L_T - 1) | \mathfrak{F}_J^T]]) \\ &\leq \mathbb{E}_{\mu_T} \left[\sup_{F \in d_1\text{-Lip}_1} (\mathbb{E}[F(L_T - 1) | \mathfrak{F}_J^T]) \right] \end{aligned}$$

But conditioning on J , the intensity is deterministic so according to Theorem 9,

$$\mathcal{T}_{d_1}(\mu_T, \nu_T) = \mathbb{E} \left[\int_0^T \left| \frac{\Psi(J_s)}{\lambda} \right| \lambda ds \exp \left[\frac{1}{2} \int_0^T \left| \frac{\Psi(J_s)}{\lambda} - 1 \right|^2 \lambda ds \right] \right],$$

which ends the proof. $\square$

One can then try to determinate the nearest Poisson process to a given MMPP. For, we seek for λ_{opt} such that the upper bound of (10) is minimal. It is clear that this minimum is directed by the exponential part of the expression. It is thus enough to minimize

$$\Phi_T(\lambda) = \int_0^T \left| \frac{\Psi(J_s)}{\lambda} - 1 \right|^2 \lambda ds$$

It is well known, that for large T , we have :

$$\Phi_T(\lambda) \underset{T \rightarrow \infty}{\sim} \lambda T \sum_{i=1}^m \left| \frac{\lambda_i}{\lambda} - 1 \right|^2 \pi(i)$$

which is minimal for $\lambda = \lambda_{opt} = \sum_{i=1}^m \lambda_i \pi(i)$. Moreover,

$$\begin{aligned} \sum_{i=1}^m \left| \frac{\lambda_i}{\lambda_{opt}} - 1 \right|^2 \lambda_{opt} \pi(i) &= \sum_{i=1}^m \frac{(\lambda_i - \lambda_{opt})^2}{\lambda_{opt}} \pi(i) \\ &= \frac{V_{opt}}{\lambda_{opt}} \end{aligned}$$

which is known in queueing theory as the burstiness of the MMPP. Finally the distance between an MMPP and the Poisson process of intensity equal the mean arrival rate of the MMPP is bounded by

$$\lambda_{opt} T \exp\left(\frac{V_{opt}}{2\lambda_{opt}} T\right).$$

In queueing theory, the choice of λ_{opt} as $\sum_{i=1}^m \lambda_i \pi(i)$ is imposed by “load” conservation: one can only compare queueing systems with the same load, i.e., the load (or traffic) is defined as the product of the mean arrival rate and of the mean service time. Our result shows that this choice is likely to be the optimal one. Moreover, we are now in position to evaluate precisely the error due to this approximation. Our bound gives a qualitative basis for the experimental rule that not only the load was important to evaluate performance of queueing system but also the so-called burstiness was to be taken into account.

REFERENCES

- [1] S. Albeverio, Yu. G. Kondratiev, and M. Röckner. Analysis and geometry on configuration spaces. *J. Funct. Anal.*, 154(2):444–500, 1998.
- [2] A. D. Barbour and T. C. Brown. Stein's method and point process approximation. *Stochastic Process. Appl.*, 43(1):9–31, 1992.
- [3] S. G. Bobkov and M. Ledoux. On modified logarithmic sobolev inequalities for bernoulli and poisson measures. *Journal of Functional Analysis*, 156(2):347–365, 1998.
- [4] N. Bouleau and F. Hirsch. *Dirichlet forms and analysis on Wiener space*, volume 14 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, 1991.
- [5] T. C. Brown and A. Xia. Stein's method and birth-death processes. *Ann. Probab.*, 29(3):1373–1403, 2001.
- [6] L. H. Y. Chen and A. Xia. Stein's method, Palm theory and Poisson process approximation. *Ann. Probab.*, 32(3B):2545–2569, 2004.
- [7] L. Decreusefond. Wasserstein distance on configuration space, 2006.
- [8] X. Fernique. Sur le théorème de Kantorovitch-Rubinstein dans les espaces polonais. In *Seminar on Probability, XV (Univ. Strasbourg, Strasbourg, 1979/1980) (French)*, volume 850 of *Lecture Notes in Math.*, pages 6–10. Springer, Berlin, 1981.
- [9] D. Feyel and A. S. Üstünel. Monge-Kantorovitch measure transportation and Monge-Ampère equation on Wiener space. *Probab. Theory Related Fields*, 128(3):347–385, 2004.
- [10] D. Feyel and A. S. Üstünel. Monge-Kantorovitch measure transportation and Monge-Ampère equation on Wiener space. *Probab. Theory Related Fields*, 128(3):347–385, 2004.
- [11] W. Fischer and K. Meier-Hellstern. The Markov-modulated Poisson process (MMPP) cookbook. *Performance Evaluation*, 18(2):149–171, 1993.
- [12] J. Jacod. *Calcul stochastique et problèmes de martingales*, volume 714 of *Lecture Notes in Mathematics*. Springer, Berlin, 1979.
- [13] M. Ledoux. Concentration of measure and logarithmic Sobolev inequalities. In *Séminaire de Probabilités, XXXIII*, volume 1709 of *Lecture Notes in Math.*, pages 120–216. Springer, Berlin, 1999.
- [14] K.S. Meier-Hellstern. The analysis of a queue arising in overflow models. *Communications, IEEE Transactions on*, 37(4):367–372, 1989.
- [15] M.F. Neuts. *Matrix-geometric solutions in stochastic models*. Dover Publications Inc., New York, 1994.
- [16] N. Privault. Chaotic and variational calculus in discrete and continuous time for the Poisson process. *Stochastics Stochastics Rep.*, 51(1-2):83–109, 1994.
- [17] S. T. Rachev and L. Rüschendorf. *Mass transportation problems. Vol. I. Probability and its Applications* (New York). Springer-Verlag, New York, 1998. Theory.
- [18] M. Röckner and A. Schied. Rademacher's theorem on configuration spaces and applications. *J. Funct. Anal.*, 169(2):325–356, 1999.
- [19] J. Ruiz de Chávez. Espaces de Fock pour les processus de Wiener et de Poisson. In *Séminaire de probabilités, XIX, 1983/84*, volume 1123 of *Lecture Notes in Math.*, pages 230–241. Springer, Berlin, 1985.
- [20] C. Villani. *Topics in optimal transportation*, volume 58 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2003.
- [21] A. Xia. A probabilistic proof of Stein's factors. *J. Appl. Probab.*, 36(1):287–290, 1999.

ENST - CNRS UMR 5141, DÉPARTEMENT INFORMATIQUE ET RÉSEAUX, 46, RUE BARRAULT, 75634 PARIS CEDEX 13, FRANCE

E-mail address: `Laurent.Decreusefond@enst.fr`

INSTITUT DE MATHÉMATIQUES DE TOULOUSE, LABORATOIRE DE STATISTIQUE ET PROBABILITÉS, UNIVERSITÉ PAUL SABATIER, 118, ROUTE DE NARBONNE, 31062 TOULOUSE CEDEX 9, FRANCE

E-mail address: `savy@math.ups-tlse.fr`